



Convex combinations of matrices – full rank characterization

Barbara Kołodziejczak, Tomasz Szulc *

*Faculty of Mathematics and Computer Science, Adam Mickiewicz University, Matejki 48/49, 60-769
Poznań, Poland*

Received 9 October 1997; accepted 29 June 1998

Submitted by V. Mehrmann

Dedicated to Ludwig Elsner on the occasion of his 60th birthday

Abstract

Let A_1, \dots, A_k be complex n -by- m matrices. Using the notion of a block P -matrix, introduced recently by Elsner and the second author, a characterization of the full rank property of all convex combinations of A_1, \dots, A_k is derived. © 1999 Elsevier Science Inc. All rights reserved.

1. Introduction

The purpose of this paper is to characterize the full rank property of all convex combinations of n -by- m complex matrices A_1, \dots, A_k .

A motivation to consider such a problem comes from the studies of solving underdetermined systems of linear algebraic equations ([1], Ch. 5.7), from the studies of solving the full rank least squares problem ([1], Ch. 5.3), and from the studies of linear systems of ordinary differential equations with regard to the topics related to controllability ([2], Appendix 2), ([3], Ch. 2.4). We also refer to the role of a full rank property in statistical analysis ([4], Ch. 4) and in research on the block-eigenvalue problem [5]. The full rank property in the case $k = 2$ has been characterized in [6]. To provide a characterization in the general

* Corresponding author. E-mail: tszulc@math.amu.edu.pl.

case we use the notion of a block P -matrix. To define it we need some notation. By $N(k)$ we will denote a fixed partition of the set $N = \{1, \dots, n\}$ into k , $1 \leq k \leq n$, pair-wise disjoint nonvoid subsets N_j of the cardinality n_j . $\mathcal{T}^{(k)}$ is then the set of all diagonal n -by- n matrices \mathbf{T} such that, for $i = 1, \dots, k$, $\mathbf{T}[N_i] = t_i \mathbf{I}_{n_i}$, where $\mathbf{T}[N_i]$ is the principal submatrix of \mathbf{T} with row and column indices in N_i , $t_i \in [0, 1]$ and \mathbf{I}_{n_i} is the n_i -by- n_i identity matrix (we set $\mathbf{I}_n = \mathbf{I}$). Now we are able to define block P -matrices.

Definition [7]. Let $N(k)$ be a partition of N . An n -by- n complex matrix \mathbf{B} is called a block P -matrix with respect to the partition $N(k)$, if for any $\mathbf{T} \in \mathcal{T}^{(k)}$

$$\det(\mathbf{TB} + (\mathbf{I} - \mathbf{T})) \neq 0.$$

2. Preliminaries

Throughout this paper it will be assumed that \mathbf{A}_i , $i = 1, \dots, k$, are n -by- m complex matrices and that $k \geq 2$. Furthermore, by Θ_k we will denote the k -by- k null matrix (we will set $\Theta_n = \Theta$).

To formulate our results we introduce some definitions. For given integer k and $\mathbf{A}_1, \dots, \mathbf{A}_k$ we define the n -by- n matrices \mathbf{A}_{ij} , $1 \leq i \leq j \leq k$, by

$$\begin{aligned} \mathbf{A}_{ii} &= \mathbf{A}_i \mathbf{A}_i^*, \quad i = 1, \dots, k, \\ \mathbf{A}_{ij} &= \mathbf{A}_i \mathbf{A}_j^* + \mathbf{A}_j \mathbf{A}_i^* - \mathbf{A}_i \mathbf{A}_i^* - \mathbf{A}_j \mathbf{A}_j^*, \quad \text{for } i < j, \end{aligned}$$

where X^* is the Hermitian adjoint of a matrix X .

Observe that, following the definitions of \mathbf{A}_{ii} and \mathbf{A}_{ij} , some manipulations will give

$$\left(\sum_{i=1}^k \alpha_i \mathbf{A}_i \right) \left(\sum_{i=1}^k \alpha_i \mathbf{A}_i \right)^* = \sum_{i=1}^k \alpha_i \mathbf{A}_{ii} + \sum_{1 \leq i < j \leq k} \alpha_i \alpha_j \mathbf{A}_{ij}, \quad (1)$$

where $\alpha_i \in [0, 1]$, $i = 1, \dots, k$, and $\sum_{i=1}^k \alpha_i = 1$.

For given integer k and \mathbf{A}_{ij} , $1 \leq i \leq j \leq k$, we define the block kn -by- kn matrices $\mathbf{B}_i = (\mathbf{B}_{sr}^i)$, $i = 1, \dots, k$, $1 \leq s, r \leq k$, by

$$\mathbf{B}_{sr}^i = \begin{cases} \mathbf{A}_{ii} & \text{if } (s, r) = (1, 1), \\ \mathbf{A}_{r-1, i} & \text{if } (s, r) \in \{(1, 2), (1, 3), \dots, (1, i)\}, \\ \mathbf{I} & \text{if } (s, r) \in \{(2, 2), (3, 3), \dots, (k, k)\}, \\ -\mathbf{I} & \text{if } (s, r) = (i+1, 1), \\ \Theta & \text{otherwise.} \end{cases}$$

So, in particular

$$\mathbf{B}_1 = \begin{bmatrix} \mathbf{A}_{11} & \mathbf{O} & \cdots & \mathbf{O} \\ -\mathbf{I} & \mathbf{I} & \cdots & \mathbf{O} \\ \cdots & \cdots & \cdots & \cdots \\ \mathbf{O} & \cdots & \mathbf{O} & \mathbf{I} \end{bmatrix}, \quad \mathbf{B}_2 = \begin{bmatrix} \mathbf{A}_{22} & \mathbf{A}_{12} & \mathbf{O} & \cdots & \mathbf{O} \\ \mathbf{O} & \mathbf{I} & \mathbf{O} & \cdots & \mathbf{O} \\ -\mathbf{I} & \mathbf{O} & \mathbf{I} & \cdots & \mathbf{O} \\ \cdots & \cdots & \cdots & \cdots & \cdots \\ \mathbf{O} & \mathbf{O} & \cdots & \mathbf{O} & \mathbf{I} \end{bmatrix},$$

$$\mathbf{B}_k = \begin{bmatrix} \mathbf{A}_{kk} & \mathbf{A}_{1k} & \cdots & \mathbf{A}_{k-1,k} \\ \mathbf{O} & \mathbf{I} & \cdots & \mathbf{O} \\ \cdots & \cdots & \cdots & \cdots \\ \mathbf{O} & \cdots & \mathbf{O} & \mathbf{I} \end{bmatrix}.$$

We close this section by recalling two results from [8] which we will need for later use.

Lemma 1. Let α_i , $i = 1, \dots, k$, be any numbers with $\sum_{i=1}^k \alpha_i = 1$. Then the following are equivalent.

- (i) $\sum_{i=1}^k \alpha_i \mathbf{A}_{ii} + \sum_{1 \leq i < j \leq k} \alpha_i \alpha_j \mathbf{A}_{ij}$ is nonsingular.
- (ii) $\sum_{i=1}^k \alpha_i \mathbf{B}_i$ is nonsingular.

Theorem 1. Let \mathbf{C}_i , $i = 1, \dots, k$, be n -by- n complex matrices. The following are equivalent.

- (i) All convex combinations of $\mathbf{C}_1, \dots, \mathbf{C}_k$ are nonsingular.
- (ii) \mathbf{C}_k is nonsingular and the $(k-1)n$ -by- $(k-1)n$ matrix

$$\begin{bmatrix} \mathbf{C}_1 \mathbf{C}_k^{-1} & (\mathbf{C}_2 - \mathbf{C}_1) \mathbf{C}_k^{-1} & (\mathbf{C}_3 - \mathbf{C}_2) \mathbf{C}_k^{-1} & \cdots & (\mathbf{C}_{k-1} - \mathbf{C}_{k-2}) \mathbf{C}_k^{-1} \\ -\mathbf{I} & \mathbf{I} & \mathbf{O} & \cdots & \mathbf{O} \\ \mathbf{O} & -\mathbf{I} & \mathbf{I} & \cdots & \mathbf{O} \\ \cdots & \cdots & \cdots & \cdots & \cdots \\ \mathbf{O} & \cdots & \mathbf{O} & -\mathbf{I} & \mathbf{I} \end{bmatrix}$$

is a block P -matrix with respect to the partition $\{M_1, \dots, M_{k-1}\}$ of $\{1, \dots, (k-1)n\}$, where $M_i = \{(i-1)n + 1, \dots, in\}$, $i = 1, \dots, k-1$.

3. Results

Theorem 2. The following are equivalent.

- (i) All convex combinations of $\mathbf{A}_1, \dots, \mathbf{A}_k$ have full row rank.
- (ii) \mathbf{A}_k has full row rank and the $(k-1)kn$ -by- $(k-1)kn$ matrix

$$\begin{bmatrix} \mathbf{B}_1 \mathbf{B}_k^{-1} & (\mathbf{B}_2 - \mathbf{B}_1) \mathbf{B}_k^{-1} & (\mathbf{B}_3 - \mathbf{B}_2) \mathbf{B}_k^{-1} & \cdots & (\mathbf{B}_{k-1} - \mathbf{B}_{k-2}) \mathbf{B}_k^{-1} \\ -\mathbf{I}_{kn} & \mathbf{I}_{kn} & \mathbf{0}_{kn} & \cdots & \mathbf{0}_{kn} \\ \mathbf{0}_{kn} & -\mathbf{I}_{kn} & \mathbf{I}_{kn} & \cdots & \mathbf{0}_{kn} \\ \cdots & \cdots & \cdots & \cdots & \cdots \\ \mathbf{0}_{kn} & \cdots & \mathbf{0}_{kn} & -\mathbf{I}_{kn} & \mathbf{I}_{kn} \end{bmatrix}$$

is a block P -matrix with respect to the partition $\{\tilde{M}_1, \dots, \tilde{M}_{k-1}\}$ of $\{1, \dots, (k-1)kn\}$, with $\tilde{M}_i = \{(i-1)kn + 1, \dots, ikn\}$, $i = 1, \dots, k-1$.

(iii) All convex combinations of $\mathbf{B}_1, \dots, \mathbf{B}_k$ are nonsingular.

Proof. We will show (ii) \Rightarrow (iii) \Rightarrow (i) \Rightarrow (ii).

First we proof that (ii) \Rightarrow (iii). As \mathbf{A}_k has full row rank hence \mathbf{A}_{kk} is nonsingular. So, \mathbf{B}_k is nonsingular and the implication in question follows by applying Theorem 1.

Now we show the implication (iii) \Rightarrow (i). From (iii), by Lemma 1 and Eq. (1), we have that for all nonnegative α_i , $i = 1, \dots, k$, with $\sum_{i=1}^k \alpha_i = 1$ the matrix

$$\left(\sum_{i=1}^k \alpha_i \mathbf{A}_i \right) \left(\sum_{i=1}^k \alpha_i \mathbf{A}_i \right)^*$$

is nonsingular. So,

$$\text{rank} \left(\left(\sum_{i=1}^k \alpha_i \mathbf{A}_i \right) \left(\sum_{i=1}^k \alpha_i \mathbf{A}_i \right)^* \right) = n.$$

But then using results on rank [9] we get

$$\text{rank} \left(\sum_{i=1}^k \alpha_i \mathbf{A}_i \right) = n$$

for all nonnegative α_i , $i = 1, \dots, k$, with $\sum_{i=1}^k \alpha_i = 1$. So, (iii) \Rightarrow (i) holds.

(i) \Rightarrow (ii): First observe that, by (i), full row rank of \mathbf{A}_k is obvious. Using again (i) and the mentioned results on rank, we get

$$\text{rank} \left(\left(\sum_{i=1}^k \alpha_i \mathbf{A}_i \right) \left(\sum_{i=1}^k \alpha_i \mathbf{A}_i \right)^* \right) = n.$$

So, the matrix

$$\left(\sum_{i=1}^k \alpha_i \mathbf{A}_i \right) \left(\sum_{i=1}^k \alpha_i \mathbf{A}_i \right)^*$$

is nonsingular. Then, by Eq. (1) and Lemma 1, all convex combinations of $\mathbf{B}_1, \dots, \mathbf{B}_k$ are nonsingular. The implication in question follows by applying Theorem 1. \square

Example 1. Let

$$\mathbf{A}_1 = \begin{bmatrix} 2 & 1.5 & 1 \\ 1 & 3 & 1 \end{bmatrix}, \quad \mathbf{A}_2 = \begin{bmatrix} 2 & 1 & 1 \\ 1 & 2.5 & 1 \end{bmatrix}, \quad \mathbf{A}_3 = \begin{bmatrix} 2 & 1 & 1 \\ 2 & 3 & 1 \end{bmatrix}.$$

As \mathbf{A}_3 is of the full row rank there exists \mathbf{B}_3^{-1} and we can form the 12-by-12 matrix

$$\mathcal{A} = \begin{bmatrix} \mathbf{B}_1 \mathbf{B}_3^{-1} & (\mathbf{B}_2 - \mathbf{B}_1) \mathbf{B}_3^{-1} \\ -\mathbf{I}_6 & \mathbf{I}_6 \end{bmatrix},$$

where

$$\mathbf{B}_1 = \begin{bmatrix} 7.25 & 7.50 & 0 & 0 & 0 & 0 \\ 7.50 & 11 & 0 & 0 & 0 & 0 \\ -1 & 0 & 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix},$$

$$\mathbf{B}_2 = \begin{bmatrix} 6 & 5.50 & -0.25 & -0.25 & 0 & 0 \\ 5.50 & 8.25 & -0.25 & -0.25 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ -1 & 0 & 0 & 0 & 1 & 0 \\ 0 & -1 & 0 & 0 & 0 & 1 \end{bmatrix},$$

$$\mathbf{B}_3^{-1} = \begin{bmatrix} 0.700 & -0.400 & 0.175 & -0.400 & 0 & -0.500 \\ -0.400 & 0.300 & -0.100 & 0.300 & 0 & 0.375 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}.$$

To check if \mathcal{A} satisfies the condition (ii) of Theorem 2 we observe that, following Theorem 3.8 from [10], a real P -matrix is a block P -matrix with respect to any partition of the index set. A direct calculation shows that all principal minors of \mathcal{A} are positive, therefore \mathcal{A} is a block P -matrix with respect to the

partition $\{\{1, \dots, 6\}, \{7, \dots, 12\}\}$ of $\{1, \dots, 12\}$. Hence, by Theorem 2, all convex combinations of A_1, A_2 and A_3 are of the full row rank.

Remark 1. In [7], another sufficient condition and some necessary conditions for a matrix to be a block P -matrix with respect to a given partition of the index set are offered. Unfortunately, a numerically verifiable characterization of this notion remains unknown.

Remark 2. In (ii) of Theorem 2 one can assume that for some i , $1 \leq i \leq k$, A_i has full row rank and the matrix in (ii) is changed accordingly. One has just to rename the A_i 's.

Remark 3. Comparing Theorem 2 with Theorem 1 it is natural to ask if the matrix in (ii) of the former result can be replaced by the $(k-1)n$ -by- $(k-1)n$ matrix

$$\begin{bmatrix} A_1 A_k^+ & (A_2 - A_1) A_k^+ & (A_3 - A_2) A_k^+ & \cdots & (A_{k-1} - A_{k-2}) A_k^+ \\ -I & I & \Theta & \cdots & \Theta \\ \Theta & -I & I & \cdots & \Theta \\ \cdots & \cdots & \cdots & \cdots & \cdots \\ \Theta & \cdots & \Theta & -I & I \end{bmatrix}, \quad (2)$$

where A_k^+ is the Moore–Penrose generalized inverse of A_k . The answer is negative as shown in the following example.

Example 2. Let $A_i = (a_j^{(i)})$, $i = 1, \dots, k-1$, be 1-by- k matrices with

$$a_j^{(i)} = \begin{cases} 1 & \text{if } j \leq i, \\ 0 & \text{otherwise} \end{cases}$$

and let $A_k = (a_j^{(k)})$ be a 1-by- k matrix with

$$a_j^{(k)} = \begin{cases} -\varepsilon & \text{if } j = 1, \\ 1 & \text{if } j = k, \\ 0 & \text{otherwise,} \end{cases}$$

where ε is a positive number.

Then it is easy to see that for any nonnegative numbers α_i , $i = 1, \dots, k$, with $\sum_{i=1}^k \alpha_i = 1$,

$$\text{rank} \left(\sum_{i=1}^k \alpha_i A_i \right) = 1.$$

On the other hand, as A_k^+ is the k -by-1 matrix of the form

$$\mathbf{A}_k^+ = \frac{1}{1 + \varepsilon^2} \begin{bmatrix} -\varepsilon \\ 0 \\ \vdots \\ 0 \\ 1 \end{bmatrix},$$

one can verify that the matrix (2) becomes the $(k-1)$ -by- $(k-1)$ lower triangular matrix $L = (l_{ij})$, $1 \leq i, j \leq k-1$ with

$$l_{ii} = \begin{cases} \frac{-\varepsilon}{1+\varepsilon^2} & \text{if } i = 1, \\ 1 & \text{otherwise.} \end{cases}$$

So, L is not a block P -matrix with respect to any partition of $\{1, \dots, k-1\}$.

Concerning Remark 3 we have the following result.

Theorem 3. Let $\text{rank}(\mathbf{A}_k) = n$ and let the $(k-1)n$ -by- $(k-1)n$ matrix (2) be a block P -matrix with respect to the partition $\{M_1, \dots, M_{k-1}\}$ of $\{1, \dots, (k-1)n\}$ with M_i , $i = 1, \dots, k-1$, defined in Theorem 1. Then all convex combinations of $\mathbf{A}_1, \dots, \mathbf{A}_k$ have full row rank.

Proof. Since $\text{rank}(\mathbf{A}_k) = n$, \mathbf{A}_{kk} is nonsingular and $\mathbf{A}_k^+ = \mathbf{A}_k^* \mathbf{A}_{kk}^{-1}$. Hence the matrix (2) becomes

$$\begin{bmatrix} \mathbf{A}_1 \mathbf{A}_k^* \mathbf{A}_{kk}^{-1} & (\mathbf{A}_2 \mathbf{A}_k^* - \mathbf{A}_1 \mathbf{A}_k^*) \mathbf{A}_{kk}^{-1} & (\mathbf{A}_3 \mathbf{A}_k^* - \mathbf{A}_2 \mathbf{A}_k^*) \mathbf{A}_{kk}^{-1} & \cdots & (\mathbf{A}_{k-1} \mathbf{A}_k^* - \mathbf{A}_{k-2} \mathbf{A}_k^*) \mathbf{A}_{kk}^{-1} \\ -\mathbf{I} & \mathbf{I} & \mathbf{0} & \cdots & \mathbf{0} \\ \mathbf{0} & -\mathbf{I} & \mathbf{I} & \cdots & \mathbf{0} \\ \cdots & \cdots & \cdots & \cdots & \cdots \\ \mathbf{0} & \cdots & \mathbf{0} & -\mathbf{I} & \mathbf{I} \end{bmatrix}.$$

Then, by Theorem 1, all convex combinations of $\mathbf{A}_1 \mathbf{A}_k^*, \dots, \mathbf{A}_k \mathbf{A}_k^*$ are nonsingular. So, we have

$$n = \text{rank} \left(\sum_{i=1}^k \alpha_i \mathbf{A}_i \mathbf{A}_k^* \right) = \text{rank} \left(\left(\sum_{i=1}^k \alpha_i \mathbf{A}_i \right) \mathbf{A}_k^* \right). \quad (3)$$

Keeping in mind that $\text{rank}(\mathbf{A}_k) = n$ from Eq. (3) it follows that $\text{rank} \left(\sum_{i=1}^k \alpha_i \mathbf{A}_i \right) = n$ for all nonnegative α_i , $i = 1, \dots, k$, with $\sum_{i=1}^k \alpha_i = 1$. So, the proof is completed. \square

Remark 4. In Theorem 3 one can assume that for some i , $1 \leq i \leq k$, $\text{rank}(\mathbf{A}_i) = n$ and the matrix (2) is changed accordingly. One has just to rename the \mathbf{A}_i 's.

Remark 5. Theorems 2 and 3 can be applied to the transposes of matrices to get conclusions on the column rank.

Acknowledgements

The authors thank two anonymous referees for valuable comments and helpful suggestions on the original version of the paper.

References

- [1] G.H. Golub, C.F. Van Loan, *Matrix Computations*, 3rd ed., Johns Hopkins University Press, Baltimore, 1996.
- [2] S. Barnett, *Matrices in Control Theory*, Van Nostrand Reinhold, London, 1971.
- [3] R.A. Horn, C.R. Johnson, *Topics in Matrix Analysis*, Cambridge University Press, Cambridge, 1991.
- [4] C.R. Rao, *Linear Statistical Inference and its Applications*, Wiley, New York, 1973.
- [5] J. Vitoria, Block eigenvalues of block compound matrices, *Linear Algebra Appl.* 47 (1982) 23–34.
- [6] T. Szulc, Rank of convex combinations of matrices, *Reliable Computing* 2 (2) (1996) 181–185.
- [7] L. Elsner, T. Szulc, Convex sets of nonsingular and block P -matrices, *Linear and Multilinear Algebra* 44 (1998) 1–12.
- [8] L. Elsner, T. Szulc, Convex combinations of matrices – nonsingularity and Schur stability characterizations, *Linear and Multilinear Algebra* (to appear).
- [9] R.A. Horn, C.R. Johnson, *Matrix Analysis*, Cambridge University Press, Cambridge, 1985.
- [10] C.R. Johnson, M. Tsatsomeros, Convex sets of nonsingular and P -matrices, *Linear and Multilinear Algebra* 38 (1995) 233–240.